

A traditional dealing with a semi-classical limit and Hopf theorem

Yanlin Yu

Abstract. This paper deals with a semi-classical limit(Theorem 1) by using traditional mathematical methods, and shows a Hopf theorem as a corollary. A formal discussing of it may be found in [7]

1 A semi-classical limit theorem

Let M be a compact, closed Riemannian manifold of dim n , and V a vector field without degenerate zeros on M . Let $\Lambda^*(M)$ be the space of differential forms on M , and

$$D = d + \delta : \Lambda^*(M) \rightarrow \Lambda^*(M)$$

be the de Rham-Hodge operator, which is an elliptic operator. Let us consider a Witten's deformation of $d + \delta$.

$$D_t = (d + \delta) + t[V^* \wedge + i(V)] : \Lambda^*(M) \rightarrow \Lambda^*(M),$$

where V^* is a 1-form dual to the vector field V , and $V^* \wedge$ means the exterior product by V^* , while $i(V)$ the interior product by V . Let

$$\square_t = D_t^2 : \Lambda^*(M) \rightarrow \Lambda^*(M),$$

and $e^{-\tau \square_t}$ be the solution operator of the heat operator $\frac{\partial}{\partial \tau} + \square_t$. It is well known that $e^{-\tau \square_t}$ is an integral operator, i.e. there exists a unique family of linear maps

$$G(\tau, q, p, t) : \Lambda_p^*(M) \rightarrow \Lambda_q^*(M)$$

such that

$$(e^{-\tau \square_t} \phi)(q) = \int_M G(\tau, q, p, t) \phi(p) dp, \quad \forall \phi.$$

Such a family of $G(\tau, q, p, t)$ is called a fundamental solution of the heat operator $\frac{\partial}{\partial \tau} + \square_t$. The fundamental solution can be determined by the following equations

$$\begin{cases} \left(\frac{\partial}{\partial \tau} + \square_t \right) G(\tau, q, p, t) = 0 \\ \lim_{\tau \rightarrow 0} \int_M G(\tau, q, p, t) \phi(p) dp = \phi(q), \quad \forall \phi, \end{cases}$$

where $\square_t$ acts on the indeterminate q . If $\square_t$ is thought as a deformation of a physical system, Witten ([3]) had considered a limit situation of $\square_t$ as $t \rightarrow \infty$, and called it a ‘‘semi-classical limit’’. By using this consideration he and [2] gave a proof of Morse inequalities. Afterwards Bismut([1]) considered a double limit

$$\lim_{s \rightarrow \infty} \lim_{\tau \rightarrow 0} \text{stre}^{-\tau \square_{\frac{s}{\tau}}}$$

to give another proof of Morse inequalities, where str means a super trace we will explain later. The first limit $\lim_{\tau \rightarrow 0}$ actually means that

$$''\tau \rightarrow 0 \quad \text{and} \quad t \rightarrow \infty \quad \text{and} \quad \tau t \quad \text{is kept as a constant.}''$$

We call this limit a semi-classical limit too, and denote it by $s - \lim$. As the semi-classical limit

$$(s - \lim) \text{stre}^{-\tau \square \frac{s}{\tau}} = (s - \lim) \int_M \text{str} G(\tau, p, p, t) dp$$

is concerned, how to understand the fundamental solution $G(\tau, q, p, t)$ is a very serious thing. We introduce $\Phi_0(\tau, t, p)$ as follows in order to replace $G(\tau, p, p, t)$ when the semi-classical limit is evaluated.

For any $p \in M$, choosing an orthonormal frame $\{E_1(p), \dots, E_n(p)\}$ at p , thus the vector $V(p)$ can be expressed as

$$V(p) = \sum_i v_i(p) E_i(p).$$

Define $v_{ij}(p)$ by

$$\nabla_{E_j(p)} V = \sum_i v_{ij}(p) E_i(p),$$

where ∇ is the Levi-Civita connection. In general the matrix $(v_{ij}(P))$ is not symmetric, we denote it by $A(p)$. Let $A(p)^*$ be the transpose of $A(p)$,

$$\theta = \theta(\tau, t, p) = 2\tau t \sqrt{A(p)A(p)^*}.$$

Define a linear map

$$\phi_0(\tau, t, p) : \Lambda_p^*(M) \rightarrow \Lambda_p^*(M)$$

by

$$\begin{aligned} \phi_0(\tau, t, p) &= \frac{1}{\sqrt{4\pi\tau}^n} \sqrt{\det\left(\frac{\theta}{\sinh\theta}\right)} \\ &\cdot \exp \left\{ -2\tau t^2 (v_1(p), \dots, v_n(p)) \frac{\cosh\theta - 1}{\theta \sinh\theta} \begin{pmatrix} v_1(p) \\ \vdots \\ v_n(p) \end{pmatrix} + \tau t \sum v_{ij}(p) E_i^+ E_j^- \right\}, \end{aligned}$$

where

$$E_j^\pm = \omega_j(p) \pm i(E_j(p)) : \Lambda_p^*(M) \rightarrow \Lambda_p^*(M),$$

$\{\omega_1(p), \dots, \omega_n(p)\}$ is the coframe dual to $\{E_1(p), \dots, E_n(p)\}$. It is easy to see that ϕ_0 does not depend on the choice of $\{E_1(p), \dots, E_n(p)\}$. In this paper we will prove the following theorem

Theorem 1 Let M be a compact closed Riemannian manifold, V a vector field without degenerated zeros. Then

$$(s - \lim) \int_M |G(\tau, p, p, t) - \phi_0(\tau, t, p)| dp = 0,$$

where we used the norm of a linear map, which is defined as usually, i.e.

$$|\psi| = \sqrt{\text{tr}(\psi\psi^*)}.$$

In order to prove theorem 1, we need to introduce a parametrix out of considerations of harmonic oscillators in §2, and by using Lemma A and Lemma B in §4 we can compare the parametrix with $G(\tau, q, p, t)$. The proof of theorem 1 will be finished in §6. In §7 we prove the Hopf theorem. The appendix, which shows an independent interest, is needed when we prove lemma A.

2 Harmonic oscillators

Let us recall Mehler formula first. It is

$$\begin{aligned}\mathcal{M}(\tau, y, x, b) = & \frac{1}{\sqrt{4\pi\tau}} \sqrt{\left(\frac{\theta}{\sinh \theta}\right)} \\ & \cdot \exp \left\{ -\frac{1}{4\tau} \frac{\theta}{\sinh \theta} [\cosh \theta \cdot (x^2 + y^2) - 2xy] \right\},\end{aligned}$$

where $\theta = 2b\tau$. The Mehler formula satisfies

$$\begin{cases} \left[\frac{\partial}{\partial \tau} - \frac{\partial^2}{\partial y^2} + b^2 y^2 \right] \mathcal{M}(\tau, y, x, b) = 0 \\ \lim_{\tau \rightarrow 0} \int_R \mathcal{M}(\tau, y, x, b) f(y) dy = f(x). \end{cases}$$

From the above equations, it is easy to see that if

$$\mathcal{M}_0(\tau, y, x, b) = \mathcal{M}(\tau, y + x, x, b),$$

then

$$\begin{cases} \left[\frac{\partial}{\partial \tau} - \frac{\partial^2}{\partial y^2} + b^2 (y + x)^2 \right] \mathcal{M}_0(\tau, y, x, b) = 0 \\ \lim_{\tau \rightarrow 0} \int_R \mathcal{M}_0(\tau, y, x, b) f(y) dy = f(0). \end{cases}$$

Again, if

$$\Phi(\tau, y, a, b) = \mathcal{M}_0(\tau, y, \frac{a}{b}, b),$$

then holds

$$\begin{cases} \left[\frac{\partial}{\partial \tau} - \frac{\partial^2}{\partial y^2} + (a + by)^2 \right] \Phi(\tau, y, a, b) = 0 \\ \lim_{\tau \rightarrow 0} \int_R \Phi(\tau, y, a, b) f(y) dy = f(0). \end{cases}$$

Let us check wheather Φ has a singularity at $b = 0$. Due to

$$\begin{aligned}\Phi(\tau, y, a, b) &= \mathcal{M}_0(\tau, y, \frac{a}{b}, b) = \mathcal{M}_0(\tau, y + \frac{a}{b}, \frac{a}{b}, b) \\ &= \frac{1}{\sqrt{4\pi\tau}} \sqrt{\frac{\theta}{\sinh \theta}} \exp \left\{ -\frac{1}{4\tau} \left[\frac{\theta \cosh \theta}{\sinh \theta} \left(\frac{a^2}{b^2} + (y + \frac{a}{b})^2 \right) - 2 \frac{\theta}{\sinh \theta} \frac{a}{b} (y + \frac{a}{b}) \right] \right\} \\ &= \frac{1}{\sqrt{4\pi\tau}} \sqrt{\frac{\theta}{\sinh \theta}} \exp \left\{ -\frac{1}{4\tau} \left[\frac{\theta \cosh \theta}{\sinh \theta} y^2 + 4a\tau \frac{\cosh \theta - 1}{\sinh \theta} y + 8a^2 \tau^2 \frac{\cosh \theta - 1}{\theta \sinh \theta} \right] \right\},\end{aligned}$$

there is no singularity at $b = 0$! From the above formula of $\Phi(\tau, y, a, b)$ it follows that

$$\begin{aligned}\Phi(\tau, y, a, b) &= \frac{1}{\sqrt{4\pi\tau}} \sqrt{\frac{\theta}{\sinh \theta}} \exp \left\{ -\frac{1}{8\tau} \frac{\theta (\cosh \theta + 1)}{\sinh \theta} y^2 \right. \\ &\quad \left. - \frac{1}{4\tau} \left[a\tau \sqrt{8 \frac{\cosh \theta - 1}{\theta \sinh \theta}} + y \sqrt{\frac{\theta (\cosh \theta - 1)}{2 \sinh \theta}} \right]^2 \right\} \\ &\leq \frac{1}{\sqrt{4\pi\tau}} \sqrt{\frac{\theta}{\sinh \theta}} \exp \left\{ -\frac{1}{8\tau} \frac{\theta (\cosh \theta + 1)}{\sinh \theta} y^2 \right\} \\ \Phi(\tau, y, a, b) &= \frac{1}{\sqrt{4\pi\tau}} \sqrt{\frac{\theta}{\sinh \theta}} \exp \left\{ -\tau a^2 \frac{\sinh \theta}{\theta \cosh \theta} - \frac{1}{4\tau} \left[\sqrt{\frac{\theta \cosh \theta}{\sinh \theta}} y + 2a\tau \frac{\cosh \theta - 1}{\sqrt{\theta \cosh \theta} \cdot \sinh \theta} \right]^2 \right\} \\ &\leq \frac{1}{\sqrt{4\pi\tau}} \sqrt{\frac{\theta}{\sinh \theta}} \exp \left\{ -\tau a^2 \frac{\sinh \theta}{\theta \cosh \theta} \right\}.\end{aligned}$$

Definition 2 For a vector $a = (a_1, \dots, a_n) \in R^n$, and an $n \times n$ matrix B , let Θ and $\Theta^\#$ be arithmetic square roots of $4\tau^2 B B^*$ and $4\tau^2 B^* B$ respectively, where B^* is the transpose of B . We define

$$\begin{aligned}\Phi(\tau, Y, a, B) &= \frac{1}{\sqrt{4\pi\tau^n}} \sqrt{\det \left(\frac{\Theta}{\sinh \Theta} \right)} \\ &\quad \exp \left\{ -\frac{1}{4\tau} Y \frac{\Theta \cosh \Theta}{\sinh \Theta} Y^* - 2\tau Y \frac{\cosh \Theta - 1}{\Theta \sinh \Theta} B a^* - 2\tau a \frac{\cosh \Theta^\# - 1}{\Theta^\# \sinh \Theta^\#} a^* \right\},\end{aligned}$$

where $Y = (y_1, \dots, y_n)$, Y^* and a^* are the transposes of Y and a respectively.

Proposition 3 $\Phi(\tau, Y, a, B)$ satisfies

$$\begin{cases} \left[\frac{\partial}{\partial \tau} - \sum_{i=1}^n \frac{\partial^2}{\partial y_i^2} + (a + YB)(a + YB)^* \right] \Phi(\tau, Y, a, B) = 0 \\ \lim_{\tau \rightarrow 0} \int_R \Phi(\tau, Y, a, B) f(Y) dY = f(0). \end{cases}$$

Proof Sometime we denote an (i, j) -element of a matrix C by C_{ij} , and $\cosh \Theta$ by $\cosh$, and $\cosh \Theta^\#$ by $\cosh^\#$. Note that $\frac{\Theta}{\tau}$ does not depend on τ , hence $\frac{\partial}{\partial \tau} \left(\frac{\Theta}{\tau} \right) = 0$. This fact helps the following computations. First

$$\begin{aligned} \frac{\partial}{\partial \tau} \Phi = & \Phi \left\{ \sqrt{\det \left(\frac{\Theta}{\tau \sinh} \right)^{-1}} \frac{\partial}{\partial \tau} \sqrt{\det \left(\frac{\Theta}{\tau \sinh} \right)} - \frac{1}{4} Y \frac{\Theta}{\tau} \frac{\partial}{\partial \tau} \left(\frac{\cosh}{\sinh} \right) Y \right. \\ & \left. - 2Y \frac{\tau}{\Theta} \frac{\partial}{\partial \tau} \left(\frac{\cosh - 1}{\sinh} \right) Ba^* - 2a \frac{\tau}{\Theta^\#} \frac{\partial}{\partial \tau} \left(\frac{\cosh^\# - 1}{\sinh^\#} \right) a^* \right\}, \end{aligned}$$

then

$$\begin{aligned} \sqrt{\det \left(\frac{\Theta}{\tau \sinh} \right)^{-1}} & \frac{\partial}{\partial \tau} \sqrt{\det \left(\frac{\Theta}{\tau \sinh} \right)} = \frac{1}{2} \left(\det \left(\frac{\Theta}{\tau \sinh} \right) \right)^{-1} \frac{\partial}{\partial \tau} \left(\det \left(\frac{\Theta}{\tau \sinh} \right) \right) \\ & = \frac{1}{2} \text{tr} \left(\left(\frac{\Theta}{\tau \sinh} \right)^{-1} \frac{\partial}{\partial \tau} \left(\frac{\Theta}{\tau \sinh} \right) \right) = \frac{1}{2} \text{tr} \left(\frac{\tau \sinh}{\Theta} \frac{\partial}{\partial \tau} \left(-\frac{\cosh}{\sinh^2} \frac{\Theta}{\tau} \right) \right) \\ & = -\frac{1}{2} \text{tr} \left(\frac{\Theta \cosh}{\tau \sinh} \right). \end{aligned}$$

And then by

$$\frac{\Theta}{\tau} \frac{\partial}{\partial \tau} \left(\frac{\cosh}{\sinh} \right) = -\frac{\Theta^2}{\tau^2 \sinh^2}, \quad \frac{\tau}{\Theta} \frac{\partial}{\partial \tau} \left(\frac{\cosh - 1}{\sinh} \right) = \frac{\cosh - 1}{\sinh^2},$$

we get

$$\frac{\partial}{\partial \tau} \Phi = \Phi \left\{ -\frac{1}{2\tau} \text{tr} \left(\frac{\Theta \cosh}{\sinh} \right) + \frac{1}{4\tau^2} Y \frac{\Theta^2}{\sinh^2} Y^* - 2Y \frac{\cosh - 1}{\sinh^2} Ba^* - 2a \frac{\cosh^\# - 1}{\sinh^{\#2}} a^* \right\}.$$

Again,

$$\begin{aligned} \frac{\partial}{\partial y_i} \Phi &= \Phi \left\{ -\frac{1}{2\tau} \left(\frac{\Theta \cosh}{\sinh} \right)_{ij} y_j - 2\tau \left(\frac{\cosh - 1}{\Theta \sinh} \right)_{ij} (Ba^*)_j \right\}, \\ \sum_i \frac{\partial^2}{\partial y_i^2} \Phi &= \Phi \left\{ -\frac{1}{2\tau} \text{tr} \left(\frac{\Theta \cosh}{\sinh} \right) + \frac{1}{4\tau^2} Y \left(\frac{\Theta \cosh}{\sinh} \right)^2 Y^* \right. \\ & \quad \left. + 2Y \frac{\cosh(\cosh - 1)}{\sinh^2} Ba^* + 4\tau^2 a B^* \left(\frac{\cosh - 1}{\Theta \sinh} \right)^2 Ba^* \right\}. \end{aligned}$$

Therefore

$$\begin{aligned} \Phi^{-1} \left(\frac{\partial}{\partial \tau} \Phi - \sum_i \frac{\partial^2}{\partial y_i^2} \Phi \right) &= -\frac{1}{4\tau^2} Y \Theta^2 Y^* - 2Y Ba^* - 2a \frac{\cosh^\# - 1}{\sinh^{\#2}} a^* \\ & \quad - 4\tau^2 a B^* \left(\frac{\cosh - 1}{\Theta \sinh} \right)^2 Ba^* \\ &= -\frac{1}{4\tau^2} Y \Theta^2 Y^* - 2Y Ba^* - 2a \frac{\cosh^\# - 1}{\sinh^{\#2}} a^* - a \left(\frac{\cosh^\# - 1}{\Theta^\# \sinh^\#} \right)^2 \Theta^{\#2} a^* \\ &= -Y B B^* Y^* - 2Y Ba^* - a a^* \\ &= -(YB + a)(B^* Y^* + a^*). \end{aligned}$$

The first half of the proposition is proved. Because the second half is easy, its proof is omitted.

Proposition 4 There hold

$$\begin{aligned} (i) \quad \Phi(\tau, Y, a, B) &\leq \frac{1}{\sqrt{4\pi\tau}} \sqrt{\det \left(\frac{\Theta}{\sinh} \right)} \exp \left\{ -\frac{1}{8\tau} Y \frac{\Theta(\cosh + 1)}{\sinh} Y^* \right\} \\ (ii) \quad \Phi(\tau, Y, a, B) &\leq \frac{1}{\sqrt{4\pi\tau}} \sqrt{\det \left(\frac{\Theta}{\sinh} \right)} \exp \left\{ -\tau a \frac{\sinh}{\Theta \cosh} a^* \right\} \end{aligned}$$

Proof The proof is similar to the case of $\Phi(\tau, y, a, b)$, which was mentioned before.

Proposition 5 Let $\Phi_0(\tau, Y, X, B) = \Phi(\tau, Y - X, XB, B)$, then

$$(i) \Phi_0(\tau, Y, a, B) = \frac{1}{\sqrt{4\pi\tau^n}} \sqrt{\det\left(\frac{\Theta}{\sinh}\right)} \exp\left\{-\frac{1}{4\tau} Y \left(\frac{\Theta \cosh}{\sinh}\right) Y^* - \frac{1}{4\tau} X \left(\frac{\Theta \cosh}{\sinh}\right) X^* + \frac{1}{2\tau} Y \left(\frac{\Theta}{\sinh}\right) X^*\right\}$$

(Mehler formula),

$$(ii) \Phi_0(\tau, Y, a, B) = \frac{1}{\sqrt{4\pi\tau^n}} \sqrt{\det\left(\frac{\Theta}{\sinh}\right)} \exp\left\{-\frac{1}{8\tau} (Y + X) \frac{\Theta(\cosh-1)}{\sinh} (Y + X)^* - \frac{1}{8\tau} (Y - X) \frac{\Theta(\cosh+1)}{\sinh} (Y - X)^*\right\}.$$

Proof We only check (i) as follows

$$\begin{aligned} & \exp\left\{-\frac{1}{4\tau} (Y - X) \frac{\Theta \cosh}{\sinh} (Y - X)^* - 2\tau (Y - X) \frac{\cosh-1}{\Theta \sinh} B (XB)^* - 2\tau (XB) \frac{\cosh^\#-1}{\Theta^\# \sinh^\#} (XB)^*\right\} \\ &= \exp\left\{-\frac{1}{4\tau} Y \frac{\Theta \cosh}{\sinh} Y^* - \frac{1}{4\tau} X \frac{\Theta \cosh}{\sinh} X^* + \frac{1}{2\tau} Y \left(\frac{\Theta \cosh}{\sinh} - 4\tau^2 \frac{\cosh-1}{\Theta \sinh} BB^*\right) X^* + \frac{1}{2\tau} X \left(4\tau^2 \frac{\cosh-1}{\Theta \sinh} BB^* - 4\tau^2 B \frac{\cosh^\#-1}{\Theta^\# \sinh^\#} B^*\right) Y^*\right\} \\ &= \exp\left\{-\frac{1}{4\tau} Y \frac{\Theta \cosh}{\sinh} Y^* - \frac{1}{4\tau} X \frac{\Theta \cosh}{\sinh} X^* + \frac{1}{2\tau} Y \frac{\Theta}{\sinh} X^*\right\}. \end{aligned}$$

Therefore (i) is true.

3 Parametrix

Choose a local orthonormal frames $\{E_1, \dots, E_n\}$ on M , let $\{\omega_1, \dots, \omega_n\}$ be the coframes dual to $\{E_1, \dots, E_n\}$, then

$$V = v_i E_i, \quad V^* = v_i \omega_i,$$

and

$$V^* \wedge + i(V) = v_i E_i^+,$$

so

$$D_t = (d + \delta) + tv_i E_i^+ = E_i^- \nabla_{E_i} + tv_i E_i^+.$$

Therefore

$$\begin{aligned} \square_t = D_t^2 &= (d + \delta)^2 + \sum_{j,k} (E_k^- \nabla_{E_k} tv_j E_j^+ + tv_j E_j^+ E_k^- \nabla_{E_k}) + t^2 \sum_{j,k} v_j v_k E_j^+ E_k^+ \\ &= (d + \delta)^2 + \sum_{j,k} tv_{jk} E_k^- E_j^+ + \sum_{j,k} tv_j (E_k^- E_j^+ + E_j^+ E_k^-) \nabla_{E_k} + t^2 \sum_j v_j^2 \\ &= (d + \delta)^2 - \sum_{j,k} tv_{jk} E_j^+ E_k^- + t^2 \sum_j v_j^2. \end{aligned}$$

(One may see [yu 6] for formulas for the multiplication table of $E_1^+, \dots, E_n^+, E_1^-, \dots, E_n^-$.)

Now for $p \in M$ in a neighbourhood of p we choose a normal coordinate system $\{y_1, \dots, y_n\}$ centering at p and an orthonormal moving frame $\{E_1, \dots, E_n\}$, which is parallel along geodesics passing through p and

$$E_i(p) = \frac{\partial}{\partial y_i} \Big|_p.$$

Of course the coordinates of p is $(0, \dots, 0)$. Suppose the coordinates of q is $(y_1, \dots, y_n) = Y$, for q near p . $\square_t$ can be written as

$$\square_t = - \sum_i \frac{\partial^2}{\partial y_i^2} + t^2 \sum_i (v_j(p) + \sum_k v_{jk}(p) y_k)^2 - \sum_{j,k} tv_{jk}(p) E_j^+ E_k^- + \dots,$$

which suggests the following definition of $H(\tau, q, p, t)$ after comparing $\square_t$ with the first equation in Proposition 3. $H(\tau, q, p, t)$ is chosen for a good approximation of the fundamental solution $G(\tau, q, p, t)$.

Definition 6 Define

$$H(\tau, q, p, t) : \Lambda_p^*(M) \rightarrow \Lambda_q^*(M)$$

by

$$H(\tau, q, p, t) = \Phi(\tau, Y, (tv_1(p), \dots, tv_n(p)), tB(p)) \exp \left\{ \sum \tau tv_{jk}(p) E_j^+ E_k^- \right\} \cdot \phi(q, p),$$

where $(B(p))_{ij} = v_{ji}(p)$, $\phi(q, p)$ is a C^∞ function, which equals 1 in a small neighbourhood of the diagonal of $M \times M$ and 0 outside a little larger neighbourhood.

The proposition 13 in §5 means that this $H(\tau, q, p, t)$ is a parametrix of $\frac{\partial}{\partial \tau} + \square_t$. Let

$$Q(\tau, q, p) = \frac{1}{\sqrt{4\pi\tau^n}} \exp \left\{ -\frac{\rho(q, p)^2}{4\tau} \right\} : \Lambda_p^*(M) \rightarrow \Lambda_q^*(M),$$

then the proposition 4 in §2 induces the following lemma immediately.

Lemma 7 For $s_0 > 0$ there exist $c_0, c_1 > 0$ such that for $\tau t < s_0$ we have

$$|H(\tau, q, p, t)| \leq c_0 Q(c_1 \tau, q, p) \exp \left\{ -\frac{\tau t^2}{c_1} v(p)^2 \right\}.$$

Proof Note that

$$|\exp \{ \tau tv_{jk}(p) E_j^+ E_k^- \}| \leq \text{const.},$$

and consider the square root of the product of right-hand sides of inequalities in the proposition 4, we get a proof easily.

Lemma 8 For $\tau_0, s_0 > 0$ there exist c_0, c_1 such that for any τ, t with $0 < \tau \leq \tau_0$ and $0 < \tau t \leq s_0$ we have

$$|(\frac{\partial}{\partial \tau} + \square_t)H(\tau, q, p, t)| \leq c_0(\sqrt{\tau}t + 1)Q(c_1 \tau, q, p) \exp \left\{ -\frac{\tau t^2}{c_1} v(p)^2 \right\}.$$

Proof First we recall some notations and facts in [5], let $H^{ij}, \Gamma_{ij}^k, R_{ijkl}$ be defined by

$$\begin{aligned} E_j &= H^{ij} \frac{\partial}{\partial y_i}, \\ \nabla_{E_i} E_j &= \Gamma_{ij}^k E_k, \\ R_{ijkl} &= -\langle (\nabla_{E_i} \nabla_{E_j} - \nabla_{E_j} \nabla_{E_i} - \nabla_{[E_i, E_j]}) E_k, E_l \rangle, \end{aligned}$$

and let (H_{ij}) be the inverse of (H^{ij}) . A corollary 8 in [5] claimed a Taylor's expansion

$$H_{ij}(y) = \delta_{ij} + \frac{1}{6} \sum_{k,l} R_{ijkl}(p) y_k y_l + \dots$$

As usual Δ_0 is defined by

$$\Delta_0 = \sum_i (\nabla_{E_i} \nabla_{E_i} - \nabla_{F_i}),$$

where $F_i = \nabla_{E_i} E_i$. Weizenbock formula reads

$$(d + \delta)^2 = -\Delta_0 + \tilde{R},$$

where

$$\tilde{R} = \frac{1}{8} \sum R_{ijkl} E_i^- E_j^- E_k^+ E_l^+ + \frac{1}{4} \sum_{i,j} R_{ijji}.$$

Thus

$$\square_t = -\Delta_0 - \sum_{jk} tv_{jk} E_j^+ E_k^- + t^2 \sum_j v_j^2 + \tilde{R}.$$

Denote

$$\exp \left\{ \sum_{jk} \tau tv_{jk}(p) E_j^+ E_k^- \right\} \phi(q, p)$$

by U , then $H = \Phi U$. By the proposition 3 in §2, Weizenbock formula and a popular formula

$$\Delta_0(\Phi U) = (\Delta_0 \Phi)U + 2 \sum_i (\nabla_{E_i} \Phi)(\nabla_{E_i} U) + \Phi(\Delta_0 U),$$

we have

$$\begin{aligned} \left(\frac{\partial}{\partial \tau} + \square_t \right)(\Phi U) = & -[(\Delta_0 - \sum_i \frac{\partial^2}{\partial y_i^2})\Phi]U - 2 \sum_i (\nabla_{E_i} \Phi)(\nabla_{E_i} U) \\ & + I_1 + I_2 + I_3 + I_4, \end{aligned}$$

where

$$\begin{aligned} I_1 &= \Phi \left(\frac{\partial}{\partial \tau} U - \sum tv_{jk} E_j^+ E_k^- U \right) \\ I_2 &= t^2 [\sum_j v_j^2 - \sum_j (v_j(p) + \sum_k v_{jk}(p) y_k)^2] \Phi U, \\ I_3 &= \tilde{R} \Phi U, \\ I_4 &= \Phi \Delta_0 U. \end{aligned}$$

Note that the lemma 7 still holds if we replace H by Φ . We write the right-hand side of the inequality of the lemma 7 as

$$c_0 Q(c_1 \tau, q, p) \exp \left\{ -\frac{\tau t^2}{c_1} v(p)^2 \right\} = \sqrt{2}^n c_0 Q(2c_1 \tau, q, p) \exp \left\{ -\frac{\tau t^2}{2c_1} v(p)^2 \right\} \xi,$$

where

$$\xi = \exp \left\{ -\frac{1}{8c_1 \tau} \rho^2(q, p) - \frac{\tau t^2}{2c_1} v(p)^2 \right\}.$$

If we prove

$$|\xi \Phi^{-1} \left(\frac{\partial}{\partial \tau} + \square_t \right)(\Phi U)| < \text{const.} (\sqrt{\tau t} + 1),$$

then

$$\begin{aligned} | \left(\frac{\partial}{\partial \tau} + \square_t \right)(\Phi U) | &= | \Phi^{-1} \left(\frac{\partial}{\partial \tau} + \square_t \right)(\Phi U) | \cdot | \Phi | \\ &\leq | \xi \Phi^{-1} \left(\frac{\partial}{\partial \tau} + \square_t \right)(\Phi U) | \sqrt{2}^n c_0 Q(2c_1 \tau, q, p) \exp \left\{ -\frac{\tau t^2}{2c_1} v(p)^2 \right\} \\ &\leq \text{const.} c_0 Q(2c_1 \tau, q, p) \exp \left\{ -\frac{\tau t^2}{2c_1} v(p)^2 \right\} (\sqrt{\tau t} + 1). \end{aligned}$$

It implies Lemma 8 is correct. So we check 6 terms in the expression of $(\frac{\partial}{\partial \tau} + \square_t)(\Phi U)$ along this way in order to enable the correctness of Lemma 8. Note that for any $m_1, m_2 > 0$

$$\left(\frac{\rho^2(q, p)}{\tau} \right)^{m_1} (\tau t^2 v(p)^2)^{m_2} \xi \leq c,$$

where the constant c depends only on m_1, m_2 . So we have

$$\begin{aligned} |\xi \Phi^{-1} I_1| &\leq \xi | \sum t(v_{jk} - v_{jk}(p)) E_j^+ E_k^- | \\ &\leq \text{const.} t |y| \xi \\ &= \text{const.} t \sqrt{\tau} \left(\frac{|y|}{\sqrt{\tau}} \right) \xi \\ &\leq \text{const.} t \sqrt{\tau}, \end{aligned}$$

$$\begin{aligned}
|\xi \Phi^{-1} I_2| &\leq t^2 |\sum (v_j(p) + v_{jk}(p) y_k + O(|y|^2))^2 - \sum (v_j(p) + v_{jk}(p) y_k)^2| \xi \\
&\leq \text{const.} t^2 (|v(p)| \cdot |y|^2 + |y|^3) \xi \\
&= \text{const.} \left[t \sqrt{\tau} (\sqrt{\tau} t |v(p)|) \left(\frac{|y|^2}{\tau} \right) + t^2 \sqrt{\tau}^3 \left(\frac{|y|}{\sqrt{\tau}} \right)^3 \right] \xi \\
&\leq \text{const.} t \sqrt{\tau},
\end{aligned}$$

where “const.” means a constant, which does not depend on τ, t, q and p . Further it is easy to see that

$$|\xi \Phi^{-1} I_3| \leq \text{const.}, \quad |\xi \Phi^{-1} I_4| \leq \text{const.}.$$

So only two terms $[-(\Delta_0 - \sum \frac{\partial^2}{\partial y_i^2}) \Phi] U$ and $[-2 \sum (\nabla_{E_i} \Phi) (\nabla_{E_i} U)]$ are left alone. From the equalities

$$\begin{aligned}
\Phi_i &= \nabla_{E_i} \Phi = H^{ij} \frac{\partial}{\partial y_j} \Phi \\
&= \Phi H^{ij} \left\{ -\frac{1}{2\tau} \left(\frac{\Theta \cosh}{\sinh} \right)_{jk} y_k - 2\tau \left(\frac{\cosh - 1}{\Theta \sinh} \right)_{jk} (Ba^*)_k \right\}
\end{aligned}$$

it follows

$$|\xi \Phi^{-1} \Phi_i| \leq \text{const.} \left(\frac{|y|}{\tau} + \tau t^2 \right) \xi.$$

Note that $\phi(q, p) = 1$ for q , which is near p , and $\exp \left\{ \sum \tau t v_{jk}(p) E_j^+ E_k^- \right\}$ does not depend on y , so due to $|\Gamma_{ij}^k| \leq \text{const.} |y|$ we have

$$|U_i| \leq \text{const.} |y|.$$

Therefore

$$|\xi \Phi^{-1} \Phi_i U_i| \leq \text{const.} \left(\frac{|y|^2}{\tau} + \tau t^2 |y| \right) \xi \leq \text{const.} (\sqrt{\tau} t + 1).$$

By using the equalities

$$\begin{aligned}
\sum_i (\Phi_{ii} - \frac{\partial^2}{\partial y_i^2} \Phi) &= \Phi (H^{ik} H^{ij} - \delta_{jk}) \left(-\frac{1}{2\tau} \right) \left(\frac{\Theta \cosh}{\sinh} \right)_{jk} \\
&\quad + \Phi H^{il} \frac{\partial H^{ij}}{\partial y_l} \left(-\frac{1}{2\tau} \right) \left(\frac{\Theta \cosh}{\sinh} \right)_{jk} y_k \\
&\quad - \Phi \Gamma_{ii}^l H^{lj} \left\{ -\frac{1}{2\tau} \left(\frac{\Theta \cosh}{\sinh} \right)_{jk} y_k - 2\tau \left(\frac{\cosh - 1}{\Theta \sinh} \right)_{jk} (Ba^*)_k \right\}
\end{aligned}$$

and

$$H^{ij} = \delta_{ij} + \frac{1}{6} \sum R_{ijkl}(p) y_k y_l + \dots$$

we get

$$|\xi \Phi^{-1} \sum_i (\Phi_{ii} - \frac{\partial^2}{\partial y_i^2} \Phi)| \leq \text{const.} \left(\frac{|y|^2}{\tau} + \tau t^2 |y| \right) \xi \leq \text{const.} (1 + \sqrt{\tau} t).$$

Therefore the lemma is proved.

4 Two lemmas

Lemma A For a fixed sufficient small $\epsilon > 0$, and for any $c_1, c_2 > 0$ with $c_1 < c_2$, there exists a constant $c = c(c_1, c_2, \epsilon)$ such that for any $q, p \in M, \nu, \tau > 0$ with $0 < \nu < \tau$, we have

$$\int_{\rho(q, z) < \epsilon} Q(c_1(\tau - \nu), q, z) Q(c_2 \nu, z, p) dz \leq c Q(c_2 \tau, q, p),$$

where

$$Q(\alpha, q, z) = \frac{1}{\sqrt{4\pi\alpha}^n} \exp \left\{ -\frac{\rho^2(q, z)}{4\alpha} \right\}.$$

Proof It is equivalent to prove

$$\int_{\rho(q, z) < \epsilon} \sqrt{\frac{\tau}{(\tau - \nu)\nu}}^n \exp \left\{ \frac{1}{4} A(\nu, \tau, q, z, p) \right\} dz \leq \text{const.},$$

where const. means a constant, which does not depend on ν, τ, q , and p , and

$$A(\nu, \tau, q, z, p) = -\frac{\rho^2(q, z)}{c_1(\tau - \nu)} - \frac{\rho^2(z, p)}{c_2\nu} + \frac{\rho^2(q, p)}{c_2\tau}.$$

Now we prove the above inequality in three separate cases.

(i) If $\rho(q, p) \geq 4\epsilon$ and $\frac{\tau - \nu}{\tau} \geq \frac{1}{2}$, then due to $\rho(q, z) < \epsilon$,

$$\rho(z, p) \geq \rho(q, p) - \rho(q, z) \geq 4\epsilon - \epsilon = 3\epsilon,$$

$$\frac{\tau - \nu}{\tau} \rho(z, p) - \frac{\nu}{\tau} \rho(q, z) \geq \frac{1}{2} \rho(z, p) - \frac{1}{2} \rho(q, z) \geq \frac{3\epsilon}{2} - \frac{\epsilon}{2} = \epsilon,$$

and thus

$$\begin{aligned} A(\nu, \tau, q, z, p) &\leq -\frac{\rho^2(q, z)}{c_1(\tau - \nu)} - \frac{\rho^2(z, p)}{c_2\nu} + \frac{\rho^2(q, p)}{c_2\tau} \\ &\leq -\frac{\rho^2(q, z)}{c_2(\tau - \nu)} - \frac{\rho^2(z, p)}{c_2\nu} + \frac{(\rho(q, z) + \rho(z, p))^2}{c_2\tau} \\ &= -\frac{\tau}{c_2(\tau - \nu)\nu} \left(\frac{\nu}{\tau} \rho(q, z) - \frac{\tau - \nu}{\tau} \rho(z, p) \right)^2 \\ &\leq -\frac{\tau}{c_2(\tau - \nu)\nu} \epsilon^2. \end{aligned}$$

Therefore

$$\sqrt{\frac{\tau}{(\tau - \nu)\nu}}^n \exp \frac{A}{4} \leq \sqrt{\frac{\tau}{(\tau - \nu)\nu}}^n \exp \left\{ -\frac{\tau}{4c_2(\tau - \nu)\nu} \epsilon^2 \right\} \leq \text{const.}$$

(ii) If $\rho(q, p) \geq 4\epsilon$ and $\frac{\tau - \nu}{\tau} < \frac{1}{2}$, then $\frac{\nu}{\tau} \geq \frac{1}{2}$ and thus

$$\begin{aligned} \sqrt{\frac{\tau}{(\tau - \nu)\nu}}^n \exp \frac{A}{4} &\leq \sqrt{\frac{2}{\tau - \nu}}^n \exp \left\{ -\left(\frac{1}{c_1} - \frac{1}{c_2} \right) \frac{\rho^2(q, z)}{4(\tau - \nu)} + \frac{1}{4} \left(-\frac{\rho^2(q, z)}{c_2(\tau - \nu)} \right. \right. \\ &\quad \left. \left. - \frac{\rho^2(z, p)}{c_2\nu} + \frac{\rho^2(q, p)}{c_2\tau} \right) \right\} \\ &\leq \sqrt{\frac{2}{\tau - \nu}}^n \exp \left\{ -\left(\frac{1}{c_1} - \frac{1}{c_2} \right) \frac{\rho^2(q, z)}{4(\tau - \nu)} - \frac{\tau}{4c_2(\tau - \nu)\nu} \left(\frac{\nu}{\tau} \rho(q, z) - \frac{\tau - \nu}{\tau} \rho(z, p) \right)^2 \right\} \\ &\leq \sqrt{\frac{2}{\tau - \nu}}^n \exp \left\{ -\left(\frac{1}{c_1} - \frac{1}{c_2} \right) \frac{\rho^2(q, z)}{4(\tau - \nu)} \right\}, \end{aligned}$$

and

$$\begin{aligned} \int_{\rho(q, z) < \epsilon} \sqrt{\frac{\tau}{(\tau - \nu)\nu}}^n \exp \left\{ \frac{A}{4} \right\} dz &\leq \int_{\rho(q, z) < \epsilon} \sqrt{\frac{2}{(\tau - \nu)}}^n \exp \left\{ -\frac{\rho^2(q, z)}{4c_0(\tau - \nu)} \right\} dz \\ &\leq \text{const.}, \end{aligned}$$

where $c_0 = (\frac{1}{c_1} - \frac{1}{c_2})^{-1}$. Therefore the lemma is true in this case.

(iii) Now we consider the case when $\rho(q, p) < 4\epsilon$. Let

$$\lambda = \frac{\nu}{\tau}, \quad \mu = \frac{\tau - \nu}{\tau}.$$

By using a reasoning in the proof of case (i),

$$\begin{aligned}\lambda\rho^2(q, z) + \mu\rho^2(z, p) - \lambda\mu\rho^2(q, p) &\geq \lambda\rho^2(q, z) + \mu\rho^2(z, p) - \lambda\mu(\rho(q, z) + \rho(z, p))^2 \\ &= (\lambda\rho(q, z) - \mu\rho(z, p))^2 \\ &\geq 0,\end{aligned}$$

we can let

$$W = \sqrt{\lambda\rho^2(q, z) + \mu\rho^2(z, p) - \lambda\mu\rho^2(q, p)}.$$

And let o be a point on the geodesic joining p and q such that

$$\frac{\rho(o, p)}{\rho(q, p)} = \lambda.$$

Without loss of generalities, we assume $n = 2$. By the last theorem in the appendix we know that for a sufficeinte small $\epsilon > 0$,

$$W^2 \geq \frac{1}{2}\rho^2(o, z).$$

Then

$$\begin{aligned}\sqrt{\frac{\tau}{(\tau-\nu)\nu}}^n \exp \frac{A}{4} &\leq \sqrt{\frac{\tau}{(\tau-\nu)\nu}}^n \exp \left\{ -\frac{\tau}{4c_2(\tau-\nu)\nu} W^2 \right\} \\ &\leq \sqrt{\frac{\tau}{(\tau-\nu)\nu}}^n \exp \left\{ -\frac{\tau}{8c_2(\tau-\nu)\nu} \rho^2(o, z) \right\}.\end{aligned}$$

Choose a geodesic coordinate system centering at o , the coordinates of a point is (ρ, θ) , where $\theta \in S^1$. Then the volume measure dz satisfies

$$dz \leq \text{const.} |\rho d\rho d\theta|.$$

And then

$$\sqrt{\frac{\tau}{(\tau-\nu)\nu}}^n \exp \frac{A}{4} dz \leq \text{const.} \sqrt{\frac{\tau}{(\tau-\nu)\nu}}^n \exp \left\{ -\frac{1}{8c_2} \frac{\tau}{(\tau-\nu)\nu} \rho^2 \right\} \rho d\rho d\theta.$$

Therefore we finish the proof of Lemma A.

Lemma B Let ϵ be small enough. For any $s_0 > 0, c > 0$ there exists a constant $h(s_0, c)$ such that for any $\tilde{c} \geq h(s_0, c)$ and $t, \tau_1, \tau_2 > 0$ with $\tau_1 t, \tau_2 t < s_0$, and $q, p \in M$ with $\rho(q, p) < \epsilon$, we have

$$\exp \left\{ -\frac{\rho^2}{4c\tau_1} - \frac{\tau_1 t^2}{c} v(p)^2 \right\} \exp \left\{ -\frac{\tau_2 t^2}{\tilde{c}} v(p)^2 \right\} \leq \exp \left\{ -\frac{\tau_2 t^2}{\tilde{c}} v(q)^2 \right\}$$

and

$$\exp \left\{ -\frac{\rho^2}{4c\tau_1} - \frac{\tau_1 t^2}{c} v(p)^2 \right\} \exp \left\{ -\frac{\tau_2 t^2}{\tilde{c}} v(q)^2 \right\} \leq \exp \left\{ -\frac{\tau_2 t^2}{\tilde{c}} v(p)^2 \right\},$$

where $\rho = \rho(q, p)$.

Proof It is equivalent to prove

$$\exp \left\{ -\frac{\rho^2}{4c\tau_1} - \frac{\tau_1 t^2}{c} v(p)^2 \right\} \exp \left\{ \frac{\tau_2 t^2}{\tilde{c}} |v(q)^2 - v(p)^2| \right\} \leq 1,$$

or

$$-\frac{\rho^2}{4c\tau_1} - \frac{\tau_1 t^2}{c} v(p)^2 + \frac{\tau_2 t^2}{\tilde{c}} |v(q)^2 - v(p)^2| \leq 0.$$

From Taylor expansion of $v(q)^2$

$$v(q)^2 = v(p)^2 + 2v(p)v'(p)\rho + \dots,$$

it follows that there exists a constant k , which does not depend on p and q , such that

$$|v(q)^2 - v(p)^2| \leq k(|v(p)|\rho + \rho^2).$$

So

$$\begin{aligned} -\frac{\rho^2}{4c\tau_1} - \frac{\tau_1 t^2}{c} v(p)^2 + \frac{\tau_2 t^2}{c} |v(q)^2 - v(p)^2| &\leq -\frac{\rho^2}{4c\tau_1} - \frac{\tau_1 t^2}{c} v(p)^2 + \frac{k\tau_2 t^2}{c} (|v(p)|\rho + \rho^2) \\ &\leq -\frac{\rho^2}{4c\tau_1} - \frac{\tau_1 t^2}{c} v(p)^2 + \frac{k\tau_2 t^2}{c} \left(\frac{2c}{t} \sqrt{\frac{v(p)^2 \tau_1 t^2}{c}} \sqrt{\frac{\rho^2}{4c\tau_1}} \right) + \frac{4kc\tau_1 \tau_2 t^2}{c} \frac{\rho^2}{4c\tau_1} \\ &\leq -\frac{\rho^2}{4c\tau_1} - \frac{\tau_1 t^2}{c} v(p)^2 + \frac{k\tau_2 t^2}{c} \left(\frac{\rho^2}{4c\tau_1} + \frac{\tau_1 t^2}{c} v(p)^2 \right) + \frac{4kc\tau_1 \tau_2 t^2}{c} \frac{\rho^2}{4c\tau_1} \\ &\leq -\frac{\rho^2}{4c\tau_1} \left(1 - \frac{ks_0 c}{c} - \frac{4ks_0^2 c}{c} \right) - \frac{\tau_1 t^2}{c} v(p)^2 \left(1 - \frac{ks_0 c}{c} \right). \end{aligned}$$

Therefore if choose $h(s_0, c) \geq (ks_0 + 4ks_0^2)c$, then we have

$$-\frac{\rho^2}{4c\tau_1} - \frac{\tau_1 t^2}{c} v(p)^2 + \frac{\tau_2 t^2}{c} |v(q)^2 - v(p)^2| \leq 0.$$

So lemma B is true.

5 Levi iteration

Definition 9 Suppose we are given $H(\tau, q, p, t)$. By the following procedure we construct $K_m(\tau, q, p, t)$, $m > 0$, and $K(\tau, q, p, t)$.

$$\begin{aligned} K_0(\tau, q, p, t) &= \left(\frac{\partial}{\partial \tau} + \square_t \right) H(\tau, q, p, t), \\ K_{m+1}(\tau, q, p, t) &= \int_0^\infty d\nu \int K_0(\tau - \nu, q, z, t) K_m(\nu, z, p, t) dz, \quad \forall m \geq 0, \\ K(\tau, q, p, t) &= \sum_{m=0}^\infty (-1)^{m+1} K_m(\tau, q, p, t). \end{aligned}$$

The above procedure is called Levi iteration.

Of course, the first question for Levi iteration is wheather the series, which defines $K(\tau, q, p, t)$, converges. We will show it does if $H(\tau, q, p, t)$ is defined by Definition 6.

Lemma 10 Choose $\phi(q, p)$ properly such that the supports of $H(\tau, q, p, t)$ and $(\frac{\partial}{\partial \tau} + \square_t)H(\tau, q, p, t)$ are contained in $\{(\tau, q, p, t) | \rho(q, p) < \epsilon\}$. Then for sufficient small ϵ and fixed $\tau_0, s_0 > 0$ there exist c_0, c_1 such that Lemma 7, Lemma 8 and the following inequalities hold.

$$\begin{aligned} |H(\tau, q, p, t)| &\leq c_0 Q(c_1 \tau, q, p) \exp \left\{ -\frac{\tau t^2}{c_1} v(q)^2 \right\}, \\ |(\frac{\partial}{\partial \tau} + \square_t)H(\tau, q, p, t)| &\leq c_0 (\sqrt{\tau} t + 1) Q(c_1 \tau, q, p) \exp \left\{ -\frac{\tau t^2}{c_1} v(q)^2 \right\}. \end{aligned}$$

Proof The lemma is trivial due to Lemma 7, Lemma 8 and Lemma B.

Lemma 11 Choose ϵ, τ_0, s_0 as in Lemma 10 and let c_0, c_1 be given in Lemma 10. Then for $\tilde{c}_1 > \text{Max}\{h(s_0, 2c_1), 2c_1, \epsilon\}$, where $h(s_0, 2c_1, \epsilon)$ is given in Lemma B, the following inequalities hold

$$|K_m(\tau, q, p, t)| \leq \left(\frac{\tilde{c}_1}{c_1} \right)^{\frac{n}{2}} c_0 (\sqrt{2}^n c_0 c)^m (\sqrt{\tau} t + 1)^{m+1} \frac{\tau^m}{m!} Q(\tilde{c}_1 \tau, q, p) \exp \left\{ -\frac{\tau t^2}{\tilde{c}_1} v(q)^2 \right\},$$

where $m \geq 0$, and $c = c(2c_1, \tilde{c}_1)$, which is given by Lemma A.

Proof Let

$$a_0 = \left(\frac{\tilde{c}_1}{c_1} \right)^{\frac{n}{2}} c_0,$$

then by Lemma 10

$$\begin{aligned}
|K_0(\tau, q, p, t)| &= |(\frac{\partial}{\partial \tau} + \square_t)H(\tau, q, p, t)| \\
&\leq c_0(\sqrt{\tau}t + 1) \frac{1}{\sqrt{4\pi c_1 \tau^n}} \exp \left\{ -\frac{\rho^2(q, p)}{4c_1 \tau} - \frac{\tau t^2}{c_1} V(q) \right\} \\
&\leq \left(\frac{\tilde{c}_1}{c_1} \right)^{\frac{n}{2}} c_0(\sqrt{\tau}t + 1) \frac{1}{\sqrt{4\pi \tilde{c}_1 \tau^n}} \exp \left\{ -\frac{\rho^2(q, p)}{4\tilde{c}_1 \tau} - \frac{\tau t^2}{\tilde{c}_1} V(q) \right\} \\
&= a_0(\sqrt{\tau}t + 1) Q(\tilde{c}_1 \tau, q, p) \exp \left\{ -\frac{\tau t^2}{\tilde{c}_1} v(q)^2 \right\}.
\end{aligned}$$

Define

$$a_{m+1} = (\sqrt{2}^n c_0 c) a_m = \left(\frac{\tilde{c}_1}{c_1} \right)^{\frac{n}{2}} c_0 (\sqrt{2}^n c_0 c)^{m+1}.$$

We are going to check the following equalities

$$|K_m(\tau, q, p, t)| \leq a_m (\sqrt{\tau}t + 1)^{m+1} \frac{\tau^m}{m!} Q(\tilde{c}_1 \tau, q, p) \exp \left\{ -\frac{\tau t^2}{\tilde{c}_1} v(q)^2 \right\},$$

by induction on m . Suppose the inequality is true for m , by using an inequality in Lemma 10 and a fact

$$\text{Support}(K_0(\tau, q, p, t)) \subset \{(\tau, q, p, t) | \rho(q, p) < \epsilon\}$$

we have

$$\begin{aligned}
|K_{m+1}| &= \left| \int_0^\tau \int_M K_0(\tau - \nu, q, z, t) K_m(\nu, z, p, t) dz d\nu \right| \\
&= \left| \int_0^\tau \int_{\rho(z, q) < \epsilon} K_0(\tau - \nu, q, z, t) K_m(\nu, z, p, t) dz d\nu \right| \\
&\leq \int_0^\tau \int_{\rho(z, q) < \epsilon} c_0 a_m (\sqrt{\tau - \nu}t + 1) (\sqrt{\nu}t + 1)^{m+1} \frac{\nu^m}{m!} \tilde{Q} dz d\nu \\
&\leq \int_0^\tau \int_{\rho(z, q) < \epsilon} c_0 a_m (\sqrt{\tau}t + 1)^{m+2} \frac{\nu^m}{m!} \tilde{Q} dz d\nu
\end{aligned}$$

where

$$\tilde{Q} = Q(c_1(\tau - \nu), q, z) \exp \left\{ -\frac{(\tau - \nu)t^2}{c_1} v(z)^2 \right\} Q(\tilde{c}_1 \nu, z, p) \exp \left\{ -\frac{\nu t^2}{\tilde{c}_1} v(z)^2 \right\}.$$

By Lemma B we have

$$\begin{aligned}
\tilde{Q} &= \sqrt{2}^n Q(2c_1(\tau - \nu), q, z) Q(\tilde{c}_1 \nu, z, p) \exp \left\{ -\frac{\rho^2(q, z)}{8(\tau - \nu)c_1} - \frac{(\tau - \nu)t^2}{2c_1} v(z)^2 \right\} \\
&\quad \exp \left\{ -\frac{(\tau - \nu)t^2}{2c_1} v(z)^2 - \frac{\nu t^2}{\tilde{c}_1} v(z)^2 \right\} \\
&= \sqrt{2}^n Q(2c_1(\tau - \nu), q, z) Q(\tilde{c}_1 \nu, z, p) \exp \left\{ -\frac{\rho^2(q, z)}{8(\tau - \nu)c_1} - \frac{(\tau - \nu)t^2}{2c_1} v(z)^2 \right\} \\
&\quad \exp \left\{ -\frac{(\tau - \nu)t^2}{2c_1} v(z)^2 - \frac{\nu t^2}{\tilde{c}_1} v(z)^2 \right\} \\
&\leq \sqrt{2}^n Q(2c_1(\tau - \nu), q, z) Q(\tilde{c}_1 \nu, z, p) \exp \left\{ -\frac{\rho^2(q, z)}{8(\tau - \nu)c_1} - \frac{(\tau - \nu)t^2}{2c_1} v(z)^2 \right\} \\
&\quad \exp \left\{ -\frac{\nu t^2}{\tilde{c}_1} v(z)^2 \right\} \\
&\leq \sqrt{2}^n Q(2c_1(\tau - \nu), q, z) Q(\tilde{c}_1 \nu, z, p) \exp \left\{ -\frac{\tau t^2}{\tilde{c}_1} v(q)^2 \right\}.
\end{aligned}$$

And by Lemma A we also have

$$\int_{\rho(z, q) < \epsilon} \tilde{Q} dz \leq \sqrt{2}^n c Q(\tilde{c}_1 \tau, q, p) \exp \left\{ -\frac{\tau t^2}{\tilde{c}_1} v(q)^2 \right\}.$$

Therefore,

$$\begin{aligned} |K_{m+1}| &\leq \int_0^\tau (c_0 c \sqrt{2}^n) a_m (\sqrt{\tau} t + 1)^{m+2} \frac{\nu^m}{m!} d\nu \cdot Q(\tilde{c}_1 \tau, q, p) \exp \left\{ -\frac{\tau t^2}{\tilde{c}_1} v(q)^2 \right\} \\ &= a_{m+1} (\sqrt{\tau} t + 1)^{m+2} \frac{\tau^{m+1}}{(m+1)!} Q(\tilde{c}_1 \tau, q, p) \exp \left\{ -\frac{\tau t^2}{\tilde{c}_1} v(q)^2 \right\}. \end{aligned}$$

The lemma is proved.

Lemma 12 There holds

$$\begin{aligned} \sum_{m=0}^\infty |K_m(\tau, q, p, t)| &\leq \left(\frac{\tilde{c}_1}{c_1} \right)^{\frac{n}{2}} c_0 (\sqrt{\tau} t + 1) \exp \left\{ \sqrt{2}^n c_0 c (\sqrt{\tau_0} s_0 + \tau_0) \right\} \\ &\quad Q(\tilde{c}_1 \tau, q, p) \exp \left\{ -\frac{\tau t^2}{\tilde{c}_1} v(q)^2 \right\}. \end{aligned}$$

Proof It is a trivial corollary of Lemma 11.

Proposition 13 There holds

$$G(\tau, q, p, t) = H(\tau, q, p, t) + \int_0^\tau d\nu \int_M H(\tau - \nu, q, z, t) K(\nu, z, p, t) dz.$$

Proof $K(\tau, q, p, t)$ is well-defined due to Lemma 12. The right-hand side of the above equality is also well-defined by using Lemma 12 and Lemma A. By a necessary routine check the right-hand side is indeed a fundamental solution of $\frac{\partial}{\partial \tau} + \square_t$. So the proposition is true.

6 Proof of Theorem 1

Due to

$$\Theta^\# = \sqrt{4\tau^2 B^* B} = \sqrt{4\tau^2 t^2 A(p) A(p)^*} = \theta$$

we have $H(\tau, p, p, t) = \phi_0(\tau, t, p)$. Then by Proposition 13, Theorem 1 is equivalent to

$$(s - \lim) \int_M dp \int_0^\tau d\nu \int_M H(\tau - \nu, p, z, t) K(\nu, z, p, t) dz = 0.$$

By lemma 10, lemma 12, and lemma B, which is used in the same way as in proving Lemma 11, we can get

$$\begin{aligned} (i) \quad |H(\tau - \nu, p, z, t) K(\nu, z, p, t)| &\leq \text{const.} \quad (\sqrt{\tau} t + 1) Q(2c_1(\tau - \nu), p, z) Q(\tilde{c}_1 \nu, z, p) \\ &\quad \exp \left\{ -\frac{\tau t^2}{\tilde{c}_1} v(p)^2 \right\}, \\ (ii) \quad |H(\tau - \nu, p, z, t) K(\nu, z, p, t)| &\leq \text{const.} \quad (\sqrt{\tau} t + 1) Q(2c_1(\tau - \nu), p, z) Q(\tilde{c}_1 \nu, z, p) \\ &\quad \exp \left\{ -\frac{\tau t^2}{\tilde{c}_1} v(z)^2 \right\}, \end{aligned}$$

where the const. is a constant, which does not depend on τ, t, q, p, z .

Let N_ϵ be an ϵ -neighbourhood of the zero set of V , and let

$$\delta = \text{Min}\{|V(p)|; p \notin N_\epsilon\}.$$

Thus if z is not in N_ϵ , then by (ii) we have

$$\begin{aligned} |H(\tau - \nu, p, z, t) K(\nu, z, p, t)| &\leq \text{const.} \quad (\sqrt{\tau} t + 1) Q(2c_1(\tau - \nu), p, z) \\ &\quad Q(\tilde{c}_1 \nu, z, p) \exp \left\{ -\frac{\tau t^2 \delta^2}{\tilde{c}_1} \right\}, \end{aligned}$$

$$\begin{aligned}
& \left| \int_M dp \int_0^\tau d\nu \int_{M-N_\epsilon} H(\tau - \nu, p, z, t) K(\nu, z, p, t) dz \right| \leq \\
& \leq \text{const.} \exp \left\{ -\frac{\tau t^2 \delta^2}{\tilde{c}_1} \right\} \int_0^\tau (\sqrt{\tau} t + 1) d\nu \\
& \quad \int \int_{\rho(p, z) < \epsilon} Q(2c_1(\tau - \nu), p, z) Q(\tilde{c}_1 \nu, z, p) dp dz \\
& \leq \text{const.} \exp \left\{ -\frac{s \delta^2}{\tilde{c}_1} t \right\} (\sqrt{\tau} s + \tau) \int_M Q(\tilde{c}_1 \tau, p, p) dp \\
& \leq \text{const.} \frac{1}{\sqrt{\tau}^\eta} \exp \left\{ -\frac{s^2 \delta^2}{\tilde{c}_1} \frac{1}{\tau} \right\},
\end{aligned}$$

which implies

$$(s - \lim) \left| \int_M dp \int_0^\tau d\nu \int_{M-N_\epsilon} H(\tau - \nu, p, z, t) K(\nu, z, p, t) dz \right| = 0.$$

Similarly, for $p \notin N_\epsilon$, we also have

$$(s - \lim) \left| \int_{M-N_\nu} dp \int_0^\tau d\nu \int_M H(\tau - \nu, p, z, t) K(\nu, z, p, t) dz \right| = 0.$$

Suppose $0 \in \text{Zero}(V)$, choose a normal coordinate system centering at 0 , and an orthonormal moving frame as in §3. Let the coordinates of p and z are $(x_1, \dots, x_n), (z_1, \dots, z_n)$ respectively. And let $Y = (y_1, \dots, y_n)$ be defined by

$$\exp_z \left(\sum y_i E_i(z) \right) = p.$$

Then

$$\begin{aligned}
H(\tau, p, z, t) &= \Phi(\tau, Y, t(v_1(z), \dots, v_n(z)), tB(z)) \\
&\quad \exp \left\{ \sum \tau t v_{jk}(z) E_j^+ E_k^- \right\} \phi(p, z),
\end{aligned}$$

and

$$|H(\tau, p, z, t)| \leq \text{const.} \Phi(\tau, Y, t(v_1(z), \dots, v_n(z)), tB(z)).$$

Note that

$$\begin{aligned}
Y &= (X - Z) + \dots, \\
(v_1(z), \dots, v_n(z)) &= (z_1, \dots, z_n) \begin{pmatrix} v_{11}(0) & \dots & v_{n1}(0) \\ \vdots & & \vdots \\ v_{1n}(0) & \dots & v_{nn}(0) \end{pmatrix} + \dots \\
&= ZB(0) + \dots
\end{aligned}$$

and

$$\begin{aligned}
\Phi_0(\tau, X - Z, tZB(0), tB(0)) &= \frac{1}{\sqrt{4\pi\tau}^\eta} \sqrt{\det \left(\frac{\Theta}{\sinh} \right)} \exp \left\{ -\frac{1}{8\tau} (X + Z) \frac{\Theta(\cosh - 1)}{\sinh} (X + Z)^* \right. \\
&\quad \left. -\frac{1}{8\tau} (X - Z) \frac{\Theta(\cosh + 1)}{\sinh} (X - Z)^* \right\}
\end{aligned}$$

Now we estimate $\Phi(\tau, Y, t(v_1(z), \dots, v_n(z)), tB(z))$ and $\Phi(\tau, X - Z, tZB(0), tB(0))$.

Lemma 14 For fixed $s > 0$, there exist $\epsilon, \eta > 0$ such that if $\rho(p, z) \leq \epsilon$, then

- (i) $\Phi(\tau, X - Z, tZB(0), tB(0)) \leq \text{const.} \frac{1}{\sqrt{4\pi\tau}^\eta} \exp \left\{ -\frac{1}{4\tau\eta} [(X + Z)(X + Z)^* + (X - Z)(X - Z)^*] \right\}.$
- (ii) $\Phi(\tau, Y, t(v_1(z), \dots, v_n(z)), tB(z)) \leq \text{const.} \frac{1}{\sqrt{4\pi\tau}^\eta} \exp \left\{ -\frac{1}{8\tau\eta} [(X + Z)(X + Z)^* + (X - Z)(X - Z)^*] \right\},$

where the const. depends on s .

Proof Denote

$$\begin{aligned} A_0 = & -\frac{1}{4\tau}(X-Z)\frac{\Theta_0 \cosh_0}{\sinh_0}(X-Z)^* \\ & -2\tau t^2(X-Z)\frac{\cosh_0-1}{\Theta_0 \sinh_0}B(0)(ZB(0))^* \\ & -2\tau t^2 ZB(0)\frac{\cosh_0^\#-1}{\Theta_0^\# \sinh_0^\#}(ZB(0))^*, \end{aligned}$$

and

$$\begin{aligned} A = & -\frac{1}{4\tau}Y\frac{\Theta \cosh}{\sinh}Y^* \\ & -2\tau t^2 Y\frac{\cosh-1}{\Theta \sinh}B(z)v(z)^* \\ & -2\tau t^2 v(z)\frac{\cosh^\#-1}{\Theta^\# \sinh^\#}v(z)^*, \end{aligned}$$

where

$$\begin{aligned} v(z) &= (v_1(z), \dots, v_n(z)), \\ \Theta &= \sqrt{4\tau^2 t^2 B(z)B(z)^*}, \\ \Theta_0 &= \sqrt{4\tau^2 t^2 B(0)B(0)^*} \\ \cosh_0 &= \cosh \Theta_0. \end{aligned}$$

It is trivial that there is a positive function $\alpha(\cdot)$ with $\lim_{\epsilon \rightarrow 0} \alpha(\epsilon) = 0$, such that

$$\begin{aligned} |A - A_0| &\leq \text{const.} \alpha(\epsilon) \left\{ \frac{1}{4\tau} |X - Z|^2 + 2\tau t^2 |X - Z| \cdot |Z| + 2\tau t^2 |Z|^2 \right\} \\ &\leq \text{const.} \alpha(\epsilon) \left\{ \frac{1}{\tau} |X - Z|^2 + \frac{1}{\tau} |X - Z| \cdot |Z| + \frac{1}{\tau} |Z|^2 \right\}. \end{aligned}$$

From

$$\begin{aligned} |Z| &\leq \frac{1}{2}(|X - Z| + |X + Z|), \\ |Z|^2 &\leq \frac{1}{4}(|X - Z| + |X + Z|)^2 \leq \frac{1}{2}(|X - Z|^2 + |X + Z|^2), \end{aligned}$$

it follows that

$$|A - A_0| \leq \text{const.} \alpha(\epsilon) \left\{ \frac{1}{\tau} |X - Z|^2 + |X + Z|^2 \right\}.$$

By using Proposition 5 we can choose $\eta > 0$ such that the following inequalities hold.

$$\begin{aligned} A_0 = & -\frac{1}{8}(X-Z)\frac{\Theta_0(\cosh_0+1)}{\sinh_0}(X-Z) \\ & -\frac{1}{8}(X+Z)\frac{\Theta_0(\cosh_0-1)}{\sinh_0}(X+Z) \\ \leq & -\frac{1}{4\tau\eta}|X-Z|^2 - \frac{1}{4\tau\eta}|X+Z|^2. \end{aligned}$$

So from

$$\frac{\Theta}{\sinh} \geq \text{const.} > 0$$

we get (i). And it is easy to see that

$$\begin{aligned} |\Phi| &\leq \text{const.} \frac{1}{\sqrt{4\pi\tau}} \exp A \leq \text{const.} \exp\{A - A_0\} \frac{1}{\sqrt{4\pi\tau}} \exp A_0 \\ &\leq \text{const.} \exp \left\{ \text{const.} \frac{\alpha(\epsilon)}{\tau} (|X - Z|^2 + |X + Z|^2) - \frac{1}{8\tau\eta} (|X - Z|^2 + |X + Z|^2) \right\} \\ &\quad \frac{1}{\sqrt{4\pi\tau}} \exp \left\{ -\frac{1}{8\tau\eta} (|X - Z|^2 + |X + Z|^2) \right\}. \end{aligned}$$

Choose ϵ small enough such that the term

$$\left\{ \text{const.} \frac{\alpha(\epsilon)}{\tau} (|X - Z|^2 + |X + Z|^2) - \frac{1}{2\tau\eta} (|X - Z|^2 + |X + Z|^2) \right\}$$

in the above inequalities is negative, thus the lemma is true.

Now let us continue to prove the theorem 1. By using Lemma 12 we have

$$\begin{aligned} |H(\tau - \nu, p, z, t) - K(\nu, z, p, t)| &\leq \text{const.} \Phi(\tau - \nu, p, z, t) (\sqrt{\nu}t + 1) Q(\tilde{c}_1\nu, q, p) \\ &\leq \text{const.} (\sqrt{\tau}t + 1) \frac{1}{\sqrt{\tau-\nu}} \frac{1}{\sqrt{\nu}} \exp \left\{ -\frac{1}{8(\tau-\nu)\eta} (X + Z)(X + Z)^* \right\} \\ &\quad \exp \left\{ -\frac{1}{8(\tau-\nu)\eta} (X - Z)(X - Z)^* \right\} \cdot \exp \left\{ -\frac{1}{4\nu\tilde{c}_1} (X - Z)(X - Z)^* \right\}. \end{aligned}$$

Let $W_1 = X + Z$, $W_2 = X - Z$, note that

$$\begin{aligned} b_0 &\equiv \int \frac{1}{\sqrt{\tau - \nu}^n} \exp \left\{ -\frac{1}{8(\tau - \nu)\eta} W_2 W_2^* \right\} \frac{1}{\sqrt{\nu}^n} \exp \left\{ -\frac{1}{4\nu\tilde{c}_1} W_2 W_2^* \right\} dW_2 \\ &\leq \int \frac{1}{\sqrt{(\tau - \nu)\nu}^n} \exp \left\{ -\frac{\tau}{4(\tau - \nu)\nu\tilde{c}_2} W_2 W_2^* \right\} dW_2 \\ &\leq \text{const.} \frac{1}{\sqrt{\tau}^n}, \end{aligned}$$

where

$$\tilde{c}_2 = \text{Max}\{2\eta, \tilde{c}_1\}.$$

Then

$$\begin{aligned} & \left| \int_0^\tau d\nu \iint_{\{\text{near } 0\}} H(\tau - \nu, p, z, t) K(\nu, z, p, t) dp dz \right| \\ & \leq \text{const.} \left| \int_0^\tau d\nu \iint_{\{\text{near } 0\}} H \cdot K dW_1 dW_2 \right| \\ & \leq \text{const.} \int_0^\tau (\sqrt{\tau}t + 1) d\nu \int b_0 \exp \left\{ -\frac{W_1^2}{8(\tau - \nu)\eta} \right\} dW_1 \\ & \leq \text{const.} \int_0^\tau (\sqrt{\tau}t + 1) d\nu \int \frac{1}{\sqrt{\tau}^n} \exp \left\{ -\frac{W_1^2}{8(\tau - \nu)\eta} \right\} dW_1 \\ & \leq \text{const.} \int_0^\tau (\sqrt{\tau}t + 1) d\nu \int \frac{1}{\sqrt{\tau}^n} \exp \left\{ -\frac{W_1^2}{4\tau\eta} \right\} dW_1 \\ & \leq \text{const.} (\sqrt{\tau}t + 1)\tau \\ & \leq \text{const.} (\sqrt{\tau}s + \tau) \xrightarrow{s\text{-lim}} 0, \end{aligned}$$

where $\{\text{near } 0\}$ means that both p and z are near to the point 0.

Suming up the above discussions we get

$$(s - \lim) \int_M dp \int_0^\tau d\nu \int_M H(\tau - \nu, p, z, t) K(\nu, z, p, t) dz = 0.$$

7 Hopf theorem

Hopf Theorem Given a Riemannian manifold M of dim n , and a vector field V without degenerate zeros, there holds

$$\chi(M) = \sum_{p \in \text{Zero}(V)} \frac{\det(v_{ij}(p))}{|\det(v_{ij}(p))|},$$

where $\chi(M)$ is the Euler number of M .

Proof For the superstructure of $\Lambda_p^*(M)$

$$\Lambda_p^*(M) = \Lambda_p^{\text{even}}(M) + \Lambda_p^{\text{odd}}(M),$$

define str of a linear map

$$L : \Lambda_p^*(M) \rightarrow \Lambda_p^*(M)$$

by

$$\text{str}(L) = \text{tr}(L|_{\Lambda_p^{\text{even}}(M)}) - \text{tr}(L|_{\Lambda_p^{\text{odd}}(M)}).$$

Due to the following inequalites

$$|a_1 + \cdots + a_n| \leq |a_1| + \cdots + |a_n| \leq \sqrt{n} \sqrt{a_1^2 + \cdots + a_n^2},$$

we have

$$|\text{str}(G(\tau, p, p, t) - \phi_0(\tau, t, p))| \leq \sqrt{n}|G(\tau, p, p, t) - \phi_0(\tau, t, p)|.$$

Thus the theorem 1 implies

$$\chi(M) = (s - \lim) \int_M \text{str} G(\tau, p, p, t) dp = (s - \lim) \int_M \text{str} \phi_0(\tau, t, p) dp.$$

Let N_ϵ be the ϵ -neighbourhood of $\text{Zero}(V)$. It is easy to see that there exists an $\delta > 0$ such that

$$\delta \leq \text{Min}\{|V(p)|; p \in M - N_\epsilon\},$$

and

$$\delta \leq \frac{\cosh \theta - 1}{\theta \sinh \theta} \quad \forall s \leq s_0$$

Then

$$|\int_{M-N_\epsilon} \text{str} \phi_0(\tau, t, p) dp| \leq \text{const.} \int_{M-N_\epsilon} \frac{1}{\sqrt{4\pi\tau}} e^{-2\tau s \delta^3} dp \xrightarrow{s-\lim} 0.$$

So

$$(s - \lim) \int_M \text{str} \phi_0(\tau, t, p) dp = (s - \lim) \int_{N_\epsilon} \text{str} \phi_0(\tau, t, p) dp.$$

Of course, $\text{Zero}(V)$ is a finite set $\{p_1, \dots, p_m\}$. For each zero point p_α , in its ϵ -neighbourhood $N_\epsilon(p_\alpha)$, choose a normal coordinate system centering at p_α , and orthonormal frames $\{E_1, \dots, E_n\}$ as before. Denote the normal coordinates of $q \in N_\epsilon(p_\alpha)$ by $(z_1, \dots, z_n)$. Let

$$\begin{aligned} W = (w_1, \dots, w_n) &= \sqrt{8}s(v_1(z), \dots, v_n(z)) \sqrt{\frac{\cosh \theta - 1}{\theta \sinh \theta}} \\ &= \sqrt{8}s((z_1, \dots, z_n)A(z)^* + \dots) \sqrt{\frac{\cosh \theta - 1}{\theta \sinh \theta}}. \end{aligned}$$

It is easy to see that

$$\frac{\partial W}{\partial Z} = \frac{\partial(w_1, \dots, w_n)}{\partial(z_1, \dots, z_n)}$$

is non-degenerate. Thus

$$\begin{aligned} (s - \lim) \int_{N_\epsilon(p_\alpha)} \text{str} \phi_0(\tau, t, q) dq &= \lim_{t \rightarrow \infty} \int_{N_\epsilon(p_\alpha)} \sqrt{\frac{1}{4\pi\tau^n}} \cdot \sqrt{\det \left(\frac{\theta}{\sinh \theta} \right)} \\ &\quad \exp\left\{-\frac{W^2}{4\tau}\right\} \exp\left\{s \sum v_{ij}(z) E_i^+ E_j^-\right\} (\det \frac{\partial W}{\partial Z})^{-1} dW \\ &= \left(\text{str} \exp\left\{s \sum v_{ij}(z) E_i^+ E_j^-\right\} (\det \frac{\partial W}{\partial Z})^{-1} \cdot \sqrt{\det \left(\frac{\theta}{\sinh \theta} \right)} \right)_{z=0}. \end{aligned}$$

From

$$\frac{\partial W}{\partial Z}|_{z=0} = \left(2sA(z)^* \sqrt{\frac{1}{\theta \sinh \theta}} \sqrt{2(\cosh \theta - 1)} \right)_{z=0},$$

it follows that

$$\left(\det \frac{\partial W}{\partial Z} \right)_{z=0}^{-1} = \left(\sqrt{\det \frac{\sinh \theta}{\theta}} \det(\sqrt{2(\cosh \theta - 1)})^{-1} \right)_{z=0}.$$

So

$$(s - \lim) \int_{N_\epsilon(p_\alpha)} \text{str} \phi_0(\tau, t, q) dq = \left(\det \sqrt{2(\cosh \theta - 1)} \right)_{z=p_\alpha}^{-1} \text{str} \exp\left\{s \sum v_{ij}(p_\alpha) E_i^+ E_j^-\right\}.$$

Let us consider everything when s goes to zero. First we have

$$\begin{aligned} [\det(\sqrt{2(\cosh \theta - 1)})]_{z=0} &= \det(\sqrt{\theta^2 + \dots})_{z=0} \\ &= (2s)^n |\det(v_{ij}(p_\alpha))| + O(s^{n+1}). \end{aligned}$$

Then let us consider the term $\text{str} \exp\{s \sum v_{ij}(p_\alpha) E_i^+ E_j^-\}$. Note that

$$\text{str}(E_{i_1}^+ \cdots E_{i_\lambda}^+ E_{j_1}^- \cdots E_{j_\mu}^-) = 0, \quad \text{for } \lambda + \mu < 2n.$$

Hence

$$\begin{aligned} \text{str}(\exp(s \sum v_{jk}(p_\alpha) E_j^+ E_k^-)) &= \frac{1}{n!} \text{str} \left(s \sum_{j,k} v_{jk}(p_\alpha) E_j^+ E_k^- \right)^n + O(s^{n+1}) \\ &= \frac{s^n}{n!} \sum \epsilon(j_1, \dots, j_n) \epsilon(k_1, \dots, k_n) v_{j_1 k_1}(p_\alpha) \cdots v_{j_n k_n}(p_\alpha) \text{str}(E_1^+ E_1^- \cdots E_n^+ E_n^-) \\ &\quad + O(s^{n+1}), \end{aligned}$$

where the sum runs over all permutations $(j_1, \dots, j_n), (k_1, \dots, k_n)$ of $(1, 2, \dots, n)$, and $\epsilon(j_1, \dots, j_n)$ is equal to 1 or -1 if the permutation $(j_1, \dots, j_n)$ is even or odd, respectively. Therefore, by the Proposition 3' in [6]

$$\text{str}(\exp(s \sum v_{jk}(p_\alpha) E_j^+ E_k^-)) = (2s)^n \cdot \det(v_{jk}(p)) + O(s^{n+1}).$$

From the above discussions we get


$$\begin{aligned} \chi(M) &= \lim_{s \rightarrow 0} (s - \lim) \int_M \text{str} G(\tau, q, q, t) dq \\ &= \lim_{s \rightarrow 0} (s - \lim) \int_M \text{str} \phi_0(\tau, t, q) dq \\ &= \sum_{p_\alpha \in \text{Zero}(V)} \lim_{s \rightarrow 0} (s - \lim) \int_{N_\epsilon(p_\alpha)} \text{str} \phi_0(\tau, t, q) dq \\ &= \sum_{p_\alpha \in \text{Zero}(V)} \lim_{s \rightarrow 0} [\det \sqrt{2(\cosh \theta - 1)}]^{-1} \text{str} \exp\{s \sum v_{ij}(p_\alpha) E_i^+ E_j^-\} \\ &= \sum_{p_\alpha \in \text{Zero}(V)} \frac{\det(v_{ij}(p_\alpha))}{|\det(v_{ij}(p_\alpha))|}. \end{aligned}$$

The theorem is proved.

8 Appendix

Let M be an oriented Riemannian manifold of dimension 2, $P : S(M) \rightarrow M$ be the tangent sphere bundle of M . Three tangent vector fields X_1, X_2, X_3 on $S(M)$ were well known for geometers (see the definitions in [4]). Let ξ^t, η^t be the integral flows on $S(M)$, which correspond to X_1, X_3 respectively. A geodesic triangle can be described by a set of parameters $\{u, t, \theta, l, \gamma, b, \alpha\}$, where $u \in S(M)$, and $\{t, \theta, l, \gamma, b, \alpha\}$ are arc lengths or angles, such that the set of parameters satisfies

$$\xi^t \eta^{\pi-\alpha} \xi^b \eta^{\pi-\gamma} \xi^l \eta^{\pi-\theta} u = u.$$



For a geodesic triangle, we denote

$$\begin{aligned} B &= P(u), & u_B &= u, \\ C &= P(\xi^l \eta^{\pi-\theta} u_B), & u_C &= \xi^l \eta^{\pi-\theta} u_B, \\ A &= P(\xi^b \eta^{\pi-\gamma} u_C), & u_A &= \xi^b \eta^{\pi-\gamma} u_C, \end{aligned}$$

and

$$\begin{aligned} [\vec{AB}] &= \begin{pmatrix} 1 & 0 \\ 0 & H(|AB|, u_B) \end{pmatrix}, \\ [\theta] &= \begin{pmatrix} -\cos \theta & -\sin \theta & 0 \\ \sin \theta & -\cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix}, \end{aligned}$$

and so on, where $|AB|$ means the length of the geodesic arc $\overline{AB}$. $H(t, u)$ is the unique solution to the following ODE

$$\begin{cases} \frac{d}{dt} X(t) = \begin{pmatrix} 0 & -k(p(t)) \\ 1 & 0 \end{pmatrix} X(t) \\ X(0) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \end{cases},$$

where $X(t)$ are 2×2 matrices, $u \in S(M)$, $p(t) = P(\xi^{-t} u)$.

Theorem(SAS trigonometry formulas) For not too big t and l , we can solve the geodesic triangle, i.e. there exist three functions

$$\begin{aligned} \alpha &= \alpha(t, \theta, l, u), \\ \gamma &= \gamma(t, \theta, l, u), \\ b &= b(t, \theta, l, u), \end{aligned}$$

such that

$$F(t, \theta, l, u) \equiv u, \quad \forall (t, \theta, l, u),$$

where

$$F(t, \theta, l, u) = \xi^t \eta^{\pi-\alpha(t, \theta, l, u)} \xi^{b(t, \theta, l, u)} \eta^{\pi-\gamma(t, \theta, l, u)} \xi^l \eta^{\pi-\theta} u.$$

Then there hold

$$\begin{aligned} (i) \quad & \begin{pmatrix} 0 & 0 & X_1(u)\alpha \\ 0 & 0 & X_2(u)\alpha \\ 0 & 0 & X_3(u)\alpha \end{pmatrix} [\vec{AB}] - \begin{pmatrix} X_1(u)b & 0 & 0 \\ X_2(u)b & 0 & 0 \\ X_3(u)b & 0 & 0 \end{pmatrix} [\alpha][\vec{AB}] \\ & + \begin{pmatrix} 0 & 0 & X_1(u)\gamma \\ 0 & 0 & X_2(u)\gamma \\ 0 & 0 & X_3(u)\gamma \end{pmatrix} [\vec{CA}][\alpha][\vec{AB}] \\ & = [\theta][\vec{BC}][\gamma][\vec{CA}][\alpha][\vec{AB}] - \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \\ (ii) \quad & \begin{pmatrix} 0 & 0 & \alpha_t \\ 0 & 0 & \alpha_\theta \\ 0 & 0 & \alpha_l \end{pmatrix} [\vec{AB}] - \begin{pmatrix} b_t & 0 & 0 \\ b_\theta & 0 & 0 \\ b_l & 0 & 0 \end{pmatrix} [\alpha][\vec{AB}] \\ & + \begin{pmatrix} 0 & 0 & \gamma_t \\ 0 & 0 & \gamma_\theta \\ 0 & 0 & \gamma_l \end{pmatrix} [\vec{CA}][\alpha][\vec{AB}] \\ & = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 0 & 0 \end{pmatrix} [\vec{BC}][\gamma][\vec{CA}][\alpha][\vec{AB}] \\ & + \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} [\gamma][\vec{CA}][\alpha][\vec{AB}] + \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \end{aligned}$$

where $\alpha_t = \frac{\partial \alpha}{\partial t}$.

Remark If M is a space form, then α, b, γ do not depend on u , hence (i) is just the last theorem in [4].

Proof Let

$$\Phi : (0, t_0] \times S^1 \times [0, l_0] \times S(M) \rightarrow S(M)$$

be the projection. From $\Phi = F$ it follows

$$\Phi_* = F_*.$$

From the trivial facts

$$\Phi_*\left(\frac{\partial}{\partial t}\right) = \Phi_*\left(\frac{\partial}{\partial \theta}\right) = \Phi_*\left(\frac{\partial}{\partial l}\right) = 0, \quad \Phi_*(X_i) = X_i,$$

we get

$$F_*\left(\frac{\partial}{\partial t}\right) = F_*\left(\frac{\partial}{\partial \theta}\right) = F_*\left(\frac{\partial}{\partial l}\right) = 0, \quad F_*(X_i) = X_i,$$

which turn out to be the desired (i) and (ii) due to concrete formulas of F_* . For example, let us compute $F_*\left(\frac{\partial}{\partial l}\right)$

$$\begin{aligned} F_*\left(\frac{\partial}{\partial l}\right) &= (\xi^t \eta^{\pi-\alpha} \xi^b \eta^{\pi-\gamma} \xi^l \eta^{\pi-\theta} u)_* \left(\frac{\partial}{\partial l}\right) \\ &= (\xi^t \eta^{\pi-\alpha} \xi^b \eta^{\pi-\gamma})_*(X_1(\xi^l \eta^{\pi-\theta} u_B)) \\ &\quad + (\xi^t \eta^{\pi-\alpha} \xi^b)_*(X_3(\eta^{\pi-\gamma} \xi^l \eta^{\pi-\theta} u_B)) \cdot (-\gamma_l) \\ &\quad + (\xi^t \eta^{\pi-\alpha})_*(X_1(\xi^b \eta^{\pi-\gamma} \xi^l \eta^{\pi-\theta} u_B)) \cdot b_l \\ &\quad + \xi_*^t(X_3(\eta^{\pi-\alpha} \xi^b \eta^{\pi-\gamma} \xi^l \eta^{\pi-\theta} u_B)) \cdot (-\alpha_l). \end{aligned}$$

By using Lemma 7 in [4] we have

$$\begin{aligned} F_*\left(\frac{\partial}{\partial l}\right) &= (1, 0, 0)[\gamma][\vec{CA}][\alpha][\vec{AB}] \\ &\quad - (0, 0, \gamma_l)[\vec{CA}][\alpha][\vec{AB}] \\ &\quad + (b_l, 0, 0)[\alpha][\vec{AB}] \\ &\quad - (0, 0, \alpha_l)[\vec{AB}]. \end{aligned}$$

So $F_*\left(\frac{\partial}{\partial l}\right) = 0$ is just the last line of the equality (ii).

Theorem there hold

- (i) $b_l = \cos \gamma$.
- (ii) $b_{ll} = -\frac{H_{11}(b, u_A)}{H_{21}(b, u_A)} \sin^2 \gamma$.
- (iii) there exist $\epsilon > 0$ such that for $0 < t, l \leq \epsilon$, $(b^2)_{ll} \geq 1$.

Proof The SAS formula shows

$$(0, 0, \alpha_l) - (b_l, 0, 0)[\alpha] + (0, 0, \gamma_l)[\vec{CA}][\alpha] = (1, 0, 0)[\gamma][\vec{CA}][\alpha],$$

i.e.

$$\begin{aligned} &(0, 0, \alpha_l) - (b_l, 0, 0) \begin{pmatrix} -\cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & -\cos \alpha & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ &\quad + (0, 0, \gamma_l) \begin{pmatrix} 1 & 0 \\ 0 & H(b, u_A) \end{pmatrix} \begin{pmatrix} -\cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & -\cos \alpha & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ &= (1, 0, 0) \begin{pmatrix} -\cos \gamma & -\sin \gamma & 0 \\ \sin \gamma & -\cos \gamma & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & H(b, u_A) \end{pmatrix} \begin{pmatrix} -\cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & -\cos \alpha & 0 \\ 0 & 0 & 1 \end{pmatrix}, \end{aligned}$$

which is

$$\begin{cases} b_l \cos \alpha + H_{21}(b, u_A) \gamma_l \sin \alpha = \cos \gamma \cos \alpha - H_{11}(b, u_A) \sin \gamma \sin \alpha \\ b_l \sin \alpha - H_{21}(b, u_A) \gamma_l \cos \alpha = \cos \gamma \sin \alpha + H_{11}(b, u_A) \sin \gamma \cos \alpha \\ \alpha_l + H_{22}(b, u_A) \gamma_l = -H_{12}(b, u_A) \sin \gamma. \end{cases}$$

So by the lemma 6 in [4] we get

$$\begin{cases} b_l &= \cos \gamma \\ \gamma_l &= -\frac{H_{11}(b, u_A)}{H_{21}(b, u_A)} \sin \gamma \\ \alpha_l &= \frac{H_{22}H_{11}}{H_{21}} \sin \gamma - H_{12} \sin \gamma \\ &= \frac{1}{H_{21}(b, u_A)} \sin \gamma \end{cases}$$

And

$$\begin{aligned} \frac{1}{2}(b^2)_l &= (b_l)^2 + b b_l = \cos^2 \gamma + (-\sin \gamma \gamma_l) \\ &= \cos^2 \gamma + \frac{b}{H_{21}} H_{11} \sin^2 \gamma = 1 + \left(\frac{b}{H_{21}} H_{11} - 1\right) \sin^2 \gamma. \end{aligned}$$

By the equation of $H(b, u_A)$ we know

$$\lim_{b \rightarrow 0} \frac{b}{H_{21}} = \lim_{b \rightarrow 0} H_{11} = 1,$$

therefore there exists $\epsilon > 0$, if $t, l \leq \epsilon$, we have

$$b \leq t + l \leq 2\epsilon,$$

so we can have

$$\left| \frac{b}{H_{21}} H_{11} - 1 \right| < \frac{1}{2}.$$

Thus

$$\begin{aligned} \left| \left(\frac{b}{H_{21}} H_{11} - 1 \right) \sin^2 \gamma \right| &< \frac{1}{2}, \\ \frac{1}{2}(b^2)_l &\geq 1 - \left| \left(\frac{b}{H_{21}} H_{11} - 1 \right) \sin^2 \gamma \right| \geq \frac{1}{2}. \end{aligned}$$

The theorem is true now.

Theorem There exists an $\epsilon > 0$, such that for any geodesic AB with $\rho(A, B) = |AB| = \epsilon$, for any $z \in M$, satisfying

$$\rho(z, A), \quad \rho(z, B) < \epsilon,$$

and for any $\lambda \in [0, 1]$, we have

$$\mu \rho(z, A)^2 + \lambda \rho(z, B)^2 - \lambda \mu \rho(A, B)^2 \geq \frac{1}{4} \rho(o, z)^2,$$

where $\mu = 1 - \lambda$, and o is a point on the geodesic AB such that

$$\frac{Ao}{AB} = \lambda.$$

Proof Let $\angle Aoz = \theta, l = \rho(o, z), \rho(A, B) = s$, and let

$$f(l) = \lambda \rho(z, B)^2 + \mu \rho(z, A)^2 - \lambda \mu l.$$

When AB and θ are fixed,

$$\frac{d\rho(A, z)}{dl} \Big|_{l=0} = \cos(\pi - \theta), \quad \frac{d\rho(B, z)}{dl} \Big|_{l=0} = \cos \theta,$$

thus

$$f(0) = \lambda\mu^2 s^2 + \mu\lambda^2 s^2 - \lambda\mu s^2 = 0,$$

and

$$\begin{aligned} f'(0) &= \lambda \frac{d\rho(z,B)^2}{dl} \Big|_{l=0} + \mu \frac{d\rho(z,A)^2}{dl} \Big|_{l=0} \\ &= \lambda\mu s \cos \theta + \mu\lambda \cos(\pi - \theta) = 0. \end{aligned}$$

Therefore

$$\begin{aligned} f(l) &= \frac{1}{2} f''(\tilde{l}) l^2 = \frac{1}{2} \left(\lambda \frac{d^2 \rho(z,B)^2}{dl^2} + \mu \frac{d^2 \rho(z,A)^2}{dl^2} \right) \Big|_{l=\tilde{l}} l^2 \\ &\geq \frac{1}{2} \left(\frac{\lambda}{2} + \frac{\mu}{2} \right) \rho^2 = \frac{\rho^2}{4}. \end{aligned}$$

So the theorem is proved.

Acknowledgement. The author is indebted to Prof H.Wu for supports to his works, and to Profs M.do Carmo, A.Rigas, Macio A.F.R. Also, the author was partially supported by the the Science Fund of the Chinese Academy of Sciences.

References

- [1] Bismut, J.-M., *The Witten complex and the degenerate Morse inequalities*, J.Diff.Geom., **23**(1986), 207-240.
- [2] Cycon, H., Froese, R., Kirsch, W. and Simon, B., *Schrodinger operator*, Springer-Verlag, (1987).
- [3] Witten, E., *Supersymmetry and Morse theory*, J.Diff.Geom., **17**(1982), 661-692.
- [4] Yu, YL., *Trigonometry(I)*, Acta Mathematica Sinica, New Series **11**1985, 135-160
- [5] Yu, YL., *Local index theorem for Dirac operator*, Acta Mathematica Sinica, New series, **3**(1987), 152-169.
- [6] Yu, YL., *Local index theorem for Signature operators*. Acta Mathematica Sinica, New series, **3**(1987), 363-372.
- [7] Yu, YL., *Semi-classical limit and Hopf theorem*, Proceedings of the international conference on manifolds and singularities, Chinese Mathematical Society–National Natural Science Foundation of China –Topology and Geometry Research Center, (1995), 113-122.

Institute of Mathematics
Academia Sinica
Beijing 100080
PR China
email address: yuyl@math03.math.ac.cn